

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VIVJOA™ safely and effectively. See full prescribing information for VIVJOA™.

VIVJOA™ (oteseconazole) capsules, for oral use
Initial U.S. Approval: 2022

INDICATIONS AND USAGE

VIVJOA™ is an azole antifungal indicated to reduce the incidence of recurrent vulvovaginal candidiasis (RVVC) in females with a history of RVVC who are NOT of reproductive potential. (1)

DOSAGE AND ADMINISTRATION

- There are two recommended VIVJOA dosage regimens: a VIVJOA-only regimen and a Fluconazole/VIVJOA regimen. Use one of these two dosage regimens. (2.1)
 - Administer VIVJOA orally with food. (2.1)
- For the *VIVJOA-only Dosage Regimen*: (2.2)
 - **On Day 1:** Administer VIVJOA 600 mg (as a single dose), then
 - **On Day 2:** Administer VIVJOA 450 mg (as a single dose), then
 - **Beginning on Day 14:** Administer VIVJOA 150 mg once a week (every 7 days) for 11 weeks (Weeks 2 through 12).
- For the *Fluconazole/VIVJOA Dosage Regimen*, prescribe fluconazole and: (2.3)
 - **On Day 1, Day 4, and Day 7:** Administer fluconazole 150 mg orally, then
 - **On Days 14 through 20:** Administer VIVJOA 150 mg once daily for 7 days, then
 - **Beginning on Day 28:** Administer VIVJOA 150 mg once a week (every 7 days) for 11 weeks (Weeks 4 through 14).

DOSAGE FORMS AND STRENGTHS

Capsules: 150 mg of oteseconazole (fluconazole is not supplied in the carton). (3)

CONTRAINDICATIONS

- Females of Reproductive Potential (4), (5.1), (8.3)
- Pregnant and Lactating women (4), (8.1), (8.2)
- Hypersensitivity to oteseconazole (4)

WARNINGS AND PRECAUTIONS

Embryo-Fetal Toxicity: Based on animal studies, VIVJOA may cause fetal harm. The drug exposure window of approximately 690 days (based on 5 times the half-life of oteseconazole) precludes adequate mitigation of the embryo-fetal toxicity risks. Advise patients that VIVJOA is contraindicated in females of reproductive potential, and in pregnant and lactating women because of potential risks to a fetus or breastfed infant. (5.1, 8.1, 8.2, 8.3)

ADVERSE REACTIONS

The most frequently reported adverse reactions (incidence > 2%) were headache and nausea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Mycovia Pharmaceuticals, Inc. at 1-855-299-0637 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

BCRP (Breast Cancer Resistance Protein) Substrates: Concomitant use of VIVJOA with BCRP substrates may increase the exposure of drugs that are BCRP substrates, which may increase the risk of adverse reactions associated with these drugs. Use the lowest possible starting dose of the BCRP substrate or consider reducing the dose of the substrate drugs and monitor for adverse reactions. (7.1)

USE IN SPECIFIC POPULATIONS

- **Renal Impairment:** Not recommended in severe renal impairment or ESRD (with or without dialysis). (8.6)
- **Hepatic Impairment:** Not recommended in moderate or severe hepatic impairment. (8.7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 4/2022

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Vulvovaginal Candidiasis
- 1.2 Usage

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage Overview and Important Administration Instructions
- 2.2 VIVJOA-only Dosage Regimen
- 2.3 Fluconazole/VIVJOA Dosage Regimen

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Effect of VIVJOA on Other Drugs

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use

- 8.6 Renal Impairment

- 8.7 Hepatic Impairment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Vulvovaginal Candidiasis

VIVJOA is indicated to reduce the incidence of recurrent vulvovaginal candidiasis (RVVC) in females with a history of RVVC who are NOT of reproductive potential [see *Warnings and Precautions (5.1)*, *Use in Specific Populations (8.3)*, and *Clinical Studies (14)*].

1.2 Usage

If specimens for fungal culture are obtained prior to therapy, antifungal therapy may be instituted before the results of the cultures are known. However, once these results become available, antifungal therapy should be adjusted accordingly.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Overview and Important Administration Instructions

There are two recommended VIVJOA dosage regimens: a VIVJOA-only regimen and a Fluconazole/VIVJOA regimen. Use one of the following two dosage regimens:

- VIVJOA-only dosage regimen [see *Dosage and Administration (2.2)*]
- Fluconazole/VIVJOA dosage regimen [see *Dosage and Administration (2.3)*].

Administer VIVJOA orally with food [see *Clinical Pharmacology (12.3)*]. Swallow the capsules whole. Do not chew, crush, dissolve, or open the capsules.

2.2 VIVJOA-only Dosage Regimen

For the VIVJOA-only dosage regimen:

- **On Day 1:** Administer VIVJOA 600 mg (as a single dose), then
- **On Day 2:** Administer VIVJOA 450 mg (as a single dose), then
- **Beginning on Day 14:** Administer VIVJOA 150 mg once a week (every 7 days) for 11 weeks (Weeks 2 through 12).

2.3 Fluconazole/VIVJOA Dosage Regimen

For the Fluconazole/VIVJOA dosage regimen, prescribe fluconazole and:

- **On Day 1, Day 4, and Day 7:** Administer fluconazole 150 mg orally, then
- **On Days 14 through 20:** Administer VIVJOA 150 mg once daily for 7 days, then
- **Beginning on Day 28:** Administer VIVJOA 150 mg once a week (every 7 days) for 11 weeks (Weeks 4 through 14).

3 DOSAGE FORMS AND STRENGTHS

VIVJOA Capsules: 150 mg of oteseconazole in lavender hard gelatin capsules imprinted with OTE 150 in black ink.

Fluconazole is not supplied in the carton.

4 CONTRAINDICATIONS

VIVJOA is contraindicated in:

- Females of reproductive potential [see *Warnings and Precautions (5.1)* and *Use in Specific Populations (8.3)*]

- Pregnant and lactating women [see *Warnings and Precautions (5.1)*, and *Use in Specific Populations (8.1, 8.2)*]
- Patients with known hypersensitivity to oteseconazole.

5 WARNINGS AND PRECAUTIONS

5.1 Embryo-Fetal Toxicity

VIVJOA is contraindicated in females of reproductive potential, and in pregnant and lactating women. Based on animal studies, VIVJOA may cause fetal harm. The drug exposure window of approximately 690 days (based on 5 times the half-life of oteseconazole) precludes adequate mitigation of the embryo-fetal toxicity risks. Ocular abnormalities were observed in the offspring of pregnant rats dosed at 7.5-mg/kg/day during organogenesis through lactation in pre and postnatal developmental studies. The observed ocular abnormalities included cataracts, opacities, exophthalmos/buphthalmos, optic nerve/retinal atrophy, lens degeneration and hemorrhage. Ocular abnormalities occurred at doses about 3.5 times the steady state clinical exposure seen with patients being treated for RVVC. Advise patients that VIVJOA is contraindicated in females of reproductive potential, and in pregnant and lactating women because of potential risks to a fetus or breastfed infant [see *Use in Specific Populations (8.1, 8.2, 8.3)*].

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of one drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

A total of 580 patients were treated with VIVJOA in three clinical trials (Trial 1, Trial 2, and Trial 3) [see *Clinical Studies (14)*]. Patients in the clinical trials were women with RVVC who received VIVJOA treatment for 12 weeks. The mean age of the patient population was 34 years (range:16-78 years), with 84% of patients aged 18-44 years and 16% of patients aged 45 years and older. Although females of reproductive potential were included in the clinical safety data, VIVJOA is contraindicated in females of reproductive potential due to the risk of embryo-fetal toxicity [see *Contraindications (4)*, *Warnings and Precautions (5.1)*, and *Use in Specific Populations (8.1, 8.3, 8.4)*].

The clinical trials population was 75% (435/580) White, 17% (96/580) Black or African American, 6% (36/580) Asian, and 2% (13/580) Other women. Fifteen percent (86/580) of all women were Hispanic/Latino. Patients enrolled in the induction and maintenance phases of the clinical trials were treated with different VIVJOA dosage regimens versus comparators [see *Clinical Studies (14)*].

The adverse reaction that led to discontinuation in 1 of 580 (0.2 %) VIVJOA-treated patients was allergic dermatitis. Overall, similar percentages of serious adverse reactions and adverse reactions leading to drug discontinuation were reported across the VIVJOA and comparator patient dosing groups.

The most frequently reported adverse reactions (incidence >2%) among VIVJOA-treated patients in Trial 1, Trial 2 and Trial 3 were headache (includes headache, migraines, sinus headaches) (7.4%) and nausea (3.6%).

Other Adverse Reactions

The following selected adverse reactions occurred in <2% of patients receiving VIVJOA in Trial 1, Trial 2 and Trial 3:

- *Laboratory investigations*: Increased blood creatine phosphokinase
- *Gastrointestinal disorders*: Dyspepsia
- *Vascular disorders*: Hot flush
- *Renal and urinary disorders*: Dysuria
- *Reproductive system and breast disorders*: Menorrhagia (includes genital hemorrhage, menorrhagia; menometrorrhagia; uterine hemorrhage, vaginal hemorrhage) metrorrhagia; vulvovaginal irritation (includes vulvovaginal burning sensation, vulvovaginal discomfort, and vulvovaginal pain)

Laboratory Findings

Elevations in Creatine Phosphokinase

Serum creatine phosphokinase (CPK) (an indirect marker of muscle injury/necrosis) elevations greater than or equal to 10 times the upper limit of normal were observed in 11 (1.9%) patients treated with VIVJOA versus 2 (0.7%) patients in the comparator groups during the VIVJOA clinical trials. The elevations were transient.

7 DRUG INTERACTIONS

7.1 Effect of VIVJOA on Other Drugs

BCRP (Breast Cancer Resistance Protein) Transporter Substrates

Oteseconazole is a BCRP inhibitor. Concomitant use of VIVJOA with BCRP substrates (e.g., rosuvastatin) may increase the exposure of BCRP substrates (e.g., rosuvastatin), which may increase the risk of adverse reactions associated with these drugs. Use the lowest possible starting dose of the BCRP substrate or consider reducing the dose of the substrate drug and monitor for adverse reactions [see *Clinical Pharmacology* (12.3)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

VIVJOA is contraindicated in females of reproductive potential and in pregnant women. Based on animal studies, VIVJOA may cause fetal harm when administered to pregnant women. In addition, the drug exposure window of approximately 690 days (based on 5 times the half-life of oteseconazole) precludes adequate mitigation of the embryo-fetal toxicity risks [see *Warnings and Precautions* (5.1) and *Clinical Pharmacology* (12.3)].

Ocular abnormalities were observed in a pre and postnatal animal study in the offspring of rats administered oteseconazole from Gestation Day 6 through Lactation Day 20 at doses about 3.5 times the recommended human dose based on AUC comparisons (see [Data](#)). The observed ocular abnormalities included cataracts, opacities, exophthalmos/buphthalmos, optic nerve/retinal atrophy, lens degeneration and hemorrhage.

There are limited human data in pregnant women who were exposed to VIVJOA during the clinical trials; these data are insufficient to exclude a potential risk of cataracts or other eye abnormalities in human infants.

Data

Animal Data

Rat and rabbit embryofetal development was assessed after oral administration of oteseconazole. There was no embryofetal toxicity or malformations at 40 mg/kg/day following administration of oteseconazole during organogenesis in pregnant rats at doses about 10 times the maximum human exposure for RVVC based on AUC comparisons. Abortions occurred in rabbits in the presence of maternal toxicity (reduced bodyweight gain with reduced food consumption) but there were no malformations at 15 mg/kg/day following administration of oteseconazole during organogenesis in pregnant rabbits about 6 times the maximum human exposure for RVVC based on AUC comparisons.

Ocular abnormalities including cataracts, opacities, exophthalmos/buphthalmos, optic nerve/retinal atrophy, lens degeneration and hemorrhage were observed in the offspring of rats administered oteseconazole from Gestation Day 6 through Lactation Day 20 at 7.5 mg/kg day (about 3.5 times the recommended human dose based on AUC comparisons). There were no effects on pregnancy or parturition in these pre and postnatal studies at any dose.

8.2 Lactation

Risk Summary

VIVJOA is contraindicated in lactating women and females of reproductive potential. There are no data on the presence of oteseconazole in human or animal milk or data on the effects of oteseconazole on milk production. There were no reported adverse effects in breastfed infants following maternal exposure to oteseconazole during lactation; however, given the limited duration of follow-up of the oteseconazole-exposed infants during the post-natal period, no conclusions can be drawn from these data [see *Warnings and Precautions (5.1)*].

Ocular abnormalities were observed in a pre and postnatal study in the offspring of rats administered oteseconazole from Gestation Day 6 through Lactation Day 20 at doses approximately 3.5 times the recommended human dose based on AUC comparisons [see *Use in Specific Populations (8.1)*]. The relationship between the observed animal findings and breastfed infants is unknown.

8.3 Females of Reproductive Potential

VIVJOA is contraindicated in females of reproductive potential based on animal findings. The drug exposure window of approximately 690 days (based on 5 times the half-life of oteseconazole) precludes adequate mitigation of the embryo-fetal toxicity risks [see *Warnings and Precautions (5.1)*, *Use in Specific Populations (8.1)* and *Clinical Pharmacology (12.3)*].

Females who are NOT of reproductive potential are defined as: persons who are biological females who are postmenopausal or have another reason for permanent infertility (e.g., tubal ligation, hysterectomy, salpingo-oophorectomy).

8.4 Pediatric Use

VIVJOA is contraindicated in females of reproductive potential. Based on animal studies, VIVJOA may cause fetal harm when administered to a pregnant woman or potential harm to the breastfed infant. The drug exposure window of approximately 690 days (based on 5 times the half-life of oteseconazole) precludes adequate mitigation of the embryo-fetal toxicity risks associated with VIVJOA use [see *Contraindications (4)*, *Warnings and Precautions (5.1)* and *Use in Specific Populations (8.1, 8.2, 8.3)* and *Clinical Pharmacology (12.3)*].

The safety and effectiveness of VIVJOA have not been established in pre-menarchal pediatric females.

8.5 Geriatric Use

Clinical studies of VIVJOA did not include sufficient numbers of patients 65 years of age and older to determine whether they respond differently from younger adult patients.

8.6 Renal Impairment

No dosage adjustment of VIVJOA is recommended in patients with mild to moderate renal impairment (i.e., estimated glomerular filtration rate (eGFR) by the modification of diet in renal disease (MDRD) equation 30-89 mL/min). Clinical studies of VIVJOA did not include sufficient numbers of patients with severe renal impairment (eGFR 15-29 mL/min) or end-stage renal disease (ESRD), defined as eGFR <15 mL/min, to determine the safety of VIVJOA in this population. Therefore, VIVJOA is not recommended for use in patients with severe renal impairment or ESRD (with or without dialysis) [see *Clinical Pharmacology* (12.3)].


8.7 Hepatic Impairment

No dosage adjustment of VIVJOA is recommended in patients with mild hepatic impairment (Child-Pugh A). There is insufficient information to determine the safety of VIVJOA in patients with moderate or severe hepatic impairment (Child-Pugh B-C). Therefore, VIVJOA is not recommended for use in patients with moderate or severe hepatic impairment [see *Clinical Pharmacology* (12.3)].

11 DESCRIPTION

VIVJOA (oteseconazole capsules) contains oteseconazole which is an oral azole antifungal agent.

The chemical name of oteseconazole is (*R*)-2-(2,4-difluorophenyl)-1,1-difluoro-3-(1*H*-tetrazol-1-yl)-1-(5-(4-(2,2,2-trifluoroethoxy)phenyl)pyridin-2-yl)propan-2-ol or 2-Pyridineethanol, α -(2,4-difluorophenyl)- β β -difluoro- α -(1*H*-tetrazol-1-ylmethyl)-5-(4-(2,2,2-trifluoroethoxy)phenyl)-, (α *R*)-. The empirical formula is C₂₃H₁₆F₇N₅O₂. The molecular weight is 527.39 g/mol. The structural formula is



Oteseconazole is a white to off-white crystalline powder and is practically insoluble in water within a pH range of 1 to 9 but is soluble in a variety of organic solvents.

Each oteseconazole capsule, for oral use, contains 150 mg oteseconazole and the following inactive ingredients: croscarmellose sodium, hydroxypropyl cellulose, lactose, magnesium stearate, silicified microcrystalline cellulose, and sodium lauryl sulfate. Capsule shell and print constituents: FD&C Blue #1, FD&C Red #3, gelatin, Opacode SW-9008/SW-9009 and titanium dioxide. Contains no ingredient made from a gluten-containing grain (wheat, barley, or rye).

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Oteseconazole is an antifungal drug [see *Microbiology* (12.4)].

12.2 Pharmacodynamics

Oteseconazole exposure-response relationships and the time course of pharmacodynamic response are unknown.

Cardiac Electrophysiology

At 5 times the maximum exposures for the recommended dose, VIVJOA does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

The AUC of oteseconazole increased approximately dose proportionally while the C_{max} increased less than dose proportionally over a dose range of 20 mg (0.13 times the lowest recommended dose) to 320 mg (0.53 times the highest recommended dose). The pharmacokinetic parameters of oteseconazole associated with the administration of the recommended dosing regimen of VIVJOA are presented in Table 1.

Table 1: Pharmacokinetic (PK) Parameters of Oteseconazole

PK Parameter^a	Mean (\pm SD)
C_{max} ($\mu\text{g/mL}$)	2.8 (1.25)
AUC _{24h} ($\text{h}\cdot\mu\text{g/mL}$)	64.2 (29.4)
C_{min} ($\mu\text{g/mL}$)	2.5 (1.19)

^a Following repeat dose administration of VIVJOA at the approved recommended dosage for RVVC at the end of treatment.

Absorption

The time to peak plasma concentrations of oteseconazole was approximately 5 to 10 hours.

Effect of Food

Administration of VIVJOA with a high-fat, high-calorie meal (800-1000 Calories; 50% fat) increased C_{max} and AUC_{0-72h} by 45% and 36%, but no significant differences were observed with a low-fat, low-calorie meal.

Distribution

The central volume of distribution of oteseconazole is approximately 423 L. Oteseconazole is 99.5-99.7% bound to plasma proteins. Animal studies indicated that oteseconazole exposures in vaginal tissue are comparable to plasma exposures.

Elimination

The median terminal half-life of oteseconazole is approximately 138 days.

Metabolism

Oteseconazole does not undergo significant metabolism.

Excretion

Following oral administration of radiolabeled oteseconazole, approximately 56% of the radiolabeled dose was recovered in feces primarily through biliary excretion and 26% was recovered in urine.

Specific Populations

There were no clinically significant differences in the pharmacokinetics of oteseconazole based on sex, race/ethnicity or mild to moderate renal impairment.

Drug Interaction Studies

BCRP substrates: Oteseconazole increased the C_{max} and AUC_{0-24h} of rosuvastatin, a BCRP substrate, by 118% and 114%, respectively.

Other Drugs: No clinically significant differences in the pharmacokinetics of the following drugs were observed when co-administered with oteseconazole: Midazolam (sensitive CYP3A4 substrate), ethinyl estradiol (CYP3A4 substrate), norethindrone (CYP3A4 substrate), or digoxin (P-gp substrate).

12.4 Microbiology

Mechanism of Action:

Oteseconazole is an azole metalloenzyme inhibitor targeting the fungal sterol, 14 α demethylase (CYP51), an enzyme that catalyzes an early step in the biosynthetic pathway of ergosterol, a sterol required for fungal cell membrane formation and integrity. Inhibition of CYP51 results in the accumulation of 14-methylated sterols, some of which are toxic to fungi. Through the inclusion of a tetrazole metal-binding group, oteseconazole has a lower affinity for human CYP enzymes.

Resistance:

The potential for increases in minimum inhibitory concentrations (MIC) to oteseconazole has been evaluated *in vitro* including specific mechanisms of resistance. Increases in oteseconazole MIC were associated with upregulation of the efflux pumps CDR1, MDR1, and the azole target, lanosterol 14-alpha-demethylase (CYP51). Against certain *Candida* spp. oteseconazole maintained meaningful *in vitro* activity against clinical isolates that were resistant to fluconazole.

Antimicrobial Activity:

The following *in vitro* data is available, but their clinical significance is unknown. Oteseconazole has been shown to be active against most isolates of the following microorganisms associated with RVVC [see *Indications and Usage (1.1)*]:

- *Candida albicans*
- *Candida glabrata*
- *Candida krusei*
- *Candida parapsilosis*
- *Candida tropicalis*
- *Candida lusitanae*
- *Candida dubliniensis*

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

There was no increase in the incidence of tumors following daily oral administration of oteseconazole to Tg.rasH2 mice for 6 months at doses of 5 mg/kg/day (males) and 15 mg/kg/day (females), which are up to 3 and 10 times, respectively, the maximum human exposure for RVVC based on AUC comparisons.

In an oral carcinogenicity study, Sprague Dawley rats were administered doses of 0.5, 1.5, or 5 mg/kg/day oteseconazole once daily for up to 90 weeks. The high dose was initially reduced from 5 to 3 mg/kg/day due to excess mortality in males and reduced body weight in females. In males, an increase in the incidence in Leydig cell adenomas of the testes and thyroid follicular cell adenomas/carcinomas (combined) were increased at ≥ 1.5 mg/kg/day (similar to the maximum human exposure for RVVC based on AUC comparisons). In females, thyroid follicular cell adenomas and carcinomas (combined) were increased at ≥ 1.5 mg/kg/day (similar to the maximum human exposure for RVVC based on AUC comparisons) and thyroid carcinomas were increased at 5 to 3 mg/kg/day

(5 times the maximum human exposure for RVVC based on AUC comparisons). The Leydig and thyroid findings are of uncertain relevance to humans.

Mutagenesis

Oteseconazole was negative in the bacterial reverse mutation assay, *in vitro* chromosomal aberration assay and micronucleus assay in rats.

Impairment of Fertility

Male rats were administered daily oral doses of 0, 0.5, 3, or 10 mg/kg/day oteseconazole beginning 42 days prior to pairing with untreated females, through the mating and post-mating period until euthanasia on Day 76 of treatment followed by a 12-week recovery period. There were no effects on reproductive and/or fertility parameters at the time of mating at 10 mg/kg/day (7 times the maximum human exposure for RVVC based on AUC comparisons). Increased incidences of abnormal sperm were observed at 3 mg/kg/day and sperm counts were reduced at 10 mg/kg/day. Although fertility was unaffected, sperm concentration remained reduced at the end of the recovery period.

Female rats were administered daily oral doses of 0, 1.5, 5, or 25 mg/kg/day oteseconazole beginning 28 days prior to cohabitation with untreated males, continuing throughout mating and through gestational day 7. Although there were no effects on estrous cyclicity, effects on reproductive and fertility parameters were observed at 25 mg/kg/day in the presence of maternal toxicity (11 times the maximum human exposure for RVVC based on AUC comparisons).

13.2 Animal Toxicology and/or Pharmacology

In an oral carcinogenicity study, Sprague Dawley rats were administered doses of 0.5, 1.5, or 5 mg/kg/day oteseconazole once daily for up to 90 weeks. The high dose was initially reduced from 5 to 3 mg/kg/day in males due to excess mortality. Incidences of hemorrhage were increased in the adrenals, brain, coagulating gland, ears, epididymides, head, heart, lung, nose, pancreas, pharynx, prostate, seminal vesicles, spinal cord, testes, thymus, and bladder of male CrI:CD[®](SD) rats (after 77 weeks of dosing at about 5 times the MRHD based on AUC comparisons). There were no increases in the incidence of hemorrhage in rats after 26 weeks at 5 mg/kg. The clinical relevance of these findings after very high doses (5 to 7 times the MRHD) for the lifetime of the rat remains unclear.

14 CLINICAL STUDIES

Overview of the Clinical Studies

A total of 656 adults and post-menarchal pediatric females with RVVC (defined as ≥ 3 episodes of vulvovaginal candidiasis (VVC) in a 12-month period) were randomized in two multicenter, multinational, double-blind, placebo-controlled trials: Trial 1 (NCT#03562156) and Trial 2 (NCT#03561701). A total of 219 adults and post-menarchal pediatric females with RVVC were randomized in a multicenter, double-blind trial [Trial 3 (NCT#03840616)]. Although females of reproductive potential were included in the clinical efficacy data, VIVJOA is contraindicated in females of reproductive potential due to the risk of embryo-fetal toxicity [see *Contraindications (4)*, *Warnings and Precautions (5.1)* and *Use in Specific Populations (8.1, 8.3, 8.4)*].

Trial 1 and Trial 2

Trial 1 and Trial 2 were both randomized, placebo-controlled trials evaluating the efficacy and safety of VIVJOA in the reduction of RVVC. Both trials consisted of two phases: an open-label induction phase and an 11-week maintenance phase. Patients received three sequential doses of 150 mg of fluconazole (every 72 hours) on Days, 1, 4 and 7 during the induction phase. Patients returned 14 days after the first dose of fluconazole and if the acute VVC episode was resolved (vulvovaginal

signs and symptoms score < 3) they were randomized (2:1) to receive either 150 mg of VIVJOA or placebo for 7 days followed by 11 weekly doses in the maintenance phase.

In Trial 1, a total of 483 patients were enrolled in the induction phase with 326 patients entering the maintenance phase with 217 patients randomized to VIVJOA and 109 patients randomized to placebo. A total of 182 patients (84%) in the VIVJOA group and 91 patients (83%) in the placebo group completed the trial. The mean age of patients was 34 years old (range 17-78 years old) with 85% of patients aged 18-44 years and 15% of patients aged 45 years and older. Patients were 72% White, 13% Black or African American, 14% Asian, and 8% were of Hispanic or Latino ethnicity.

In Trial 2, a total of 425 patients were enrolled into the induction phase with 330 patients entering the maintenance phase with 220 subjects randomized to VIVJOA and 110 patients randomized to placebo. A total of 191 patients (87%) in the VIVJOA group and 91 patients (83%) in the placebo group completed the trial. The mean age of patients was 34 years old (range 18-73 years old) with 85% of patients aged 18-44 years and 15% of patients aged 45 years and older. Patients were 89% White, 10% Black or African American and 15% were of Hispanic or Latino ethnicity.

For both Trial 1 and Trial 2, efficacy was assessed by the proportion of patients with ≥ 1 culture-verified acute VVC episode (positive fungal culture for *Candida* species associated with a clinical signs and symptoms score of ≥ 3) during the Maintenance Phase through Week 48. Evaluation of clinical signs and symptoms included erythema (redness), edema (swelling), excoriation (skin picking), itching, burning and irritation. Since treatment for acute VVC was allowed to be provided to a patient if it was deemed to be clinically needed when the patient had a signs and symptoms score ≥ 3 and a positive KOH test, the proportion of patients with ≥ 1 culture-verified acute VVC episode or who took medication known to treat VVC during the Maintenance Phase through Week 48 is also presented.

VIVJOA was superior to placebo with reference to the proportion of patients with ≥ 1 culture-verified acute VVC episode through Week 48 (Table 2) or the proportion of patients with ≥ 1 culture-verified acute VVC episode or who took medication known to treat VVC during the Maintenance Phase through Week 48. For both Trial 1 and Trial 2, the average percentage of patients was lower in the VIVJOA groups compared with the placebo group (Table 2).

Table 2: Trial 1 and 2 Efficacy Endpoints: ITT Population

	Trial 1		Trial 2	
	VIVJOA (N=217)	Placebo (N=109)	VIVJOA (N=218)	Placebo (N=108)
Proportion of Patients with ≥ 1 Culture-verified Acute VVC Episode (Day 1 through Week 48) ^a	6.7%	42.8%	3.9%	39.4%
Treatment Difference p-value ^b	<0.001		<0.001	
Proportion of Patients with ≥ 1 Culture-verified Acute VVC Episode or received VVC medication (Day 1 through Week 48) ^a	27.3%	50.8%	21.3%	49.7%
Treatment Difference p-value ^b	<0.001		<0.001	

Abbreviations: ITT=Intent-to-Treat (Population); VVC=vulvovaginal candidiasis.

a. Average %. Missing values were imputed with multiple imputation using the following auxiliary information: region, treatment, Baseline body mass index, Baseline age, ethnicity, and visit.

b. The p-value was obtained using a Chi-square test comparing VIVJOA with placebo.

Trial 3

Trial 3 was a randomized, double-blind trial evaluating the efficacy and safety of VIVJOA versus fluconazole and placebo in adults and post-menarchal pediatric females with RVVC. The trial consisted of two phases: induction and maintenance.

During the induction phase, patients received 1050 mg of VIVJOA over two days (600 mg [4x150mg] on Day 1 and 450 mg [3x150mg] on Day 2) or three sequential doses of 150 mg of fluconazole (every 72 hours) on Days, 1, 4 and 7. Patients returned 14 days after the first dose and moved to the maintenance phase if the acute VVC episode was resolved. During the maintenance phase, patients received 150 mg VIVJOA weekly or placebo weekly for 11 weeks.

A total of 219 patients were randomized (2:1) into the induction phase: 147 to VIVJOA and 72 to fluconazole/placebo. One patient in the VIVJOA group did not receive drug therefore 146 patients received VIVJOA. A total of 112 patients (76%) in the VIVJOA group and 55 patients (76%) in the fluconazole/placebo group completed the trial.

The mean age of patients was 35 years (range 16-78) with 80% of patients aged 18-44 years and 19% of patients aged 45 years and older. Patients were 59% White, 34% Black or African American, 1% Asian and 26% were of Hispanic or Latino ethnicity. The trial was conducted completely in the United States.

Efficacy was assessed by the proportion of patients with ≥ 1 culture verified acute VVC episode during the maintenance phase (post-randomization through Week 50) or who failed clearing their infection during the induction phase. A recurring acute VVC episode was defined as a positive culture for *Candida* species and a clinical signs and symptoms score of ≥ 3 . Evaluation of clinical signs and symptoms included erythema (redness), edema (swelling), excoriation (skin picking), itching, burning and irritation. Additionally, the proportion of patients with ≥ 1 culture verified acute VVC episode or who took medication known to treat VVC during the maintenance phase (post-randomization through Week 50) or who failed clearing their infection during the induction phase is presented.

VIVJOA was superior to fluconazole/placebo in the proportion of patients with ≥ 1 culture-verified recurring acute VVC episode during the maintenance phase (post randomization through Week 50) or failed clearing their infection during the induction phase and the proportion of patients with ≥ 1 culture-verified recurring acute VVC episode or took VVC medication known to treat VVC during the maintenance phase (post randomization through Week 50) or who failed clearing their infection during the induction phase. The average percentage of patients was lower in the VIVJOA group compared with the fluconazole/placebo group ([Table 3](#)).

Table 3: Trial 3 Efficacy Endpoints ITT population

	VIVJOA (N=147)	Fluconazole/Placebo (N=72)	Treatment Difference p-value^b
Proportion of Patients with ≥1 Culture-verified Acute VVC Episode through Week 50 or Unresolved VVC Episode During the Induction Phase ^a	10.3%	42.9%	<0.001
Proportion of Patients with ≥1 Culture-verified Acute VVC Episode or took VVC medication through Week 50 or Unresolved VVC Episode During the Induction Phase ^a	43.5%	59.0%	0.039

Abbreviations: ITT=Intent-to-Treat (Population); VVC=vulvovaginal candidiasis

a. Average %, Missing values were imputed with multiple imputation using the following auxiliary information: treatment, baseline body mass index, baseline age, ethnicity, and visit.

b. The p-value was obtained using a Chi-square test comparing VIVJOA with fluconazole/placebo.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

VIVJOA (oteseconazole capsules) are supplied as lavender hard gelatin capsules. Printed black “OTE 150” on the capsule and contain 150 mg oteseconazole. They are available in an 18-count (NDC 74695-823-18) blister package within a child resistant wallet. There will be one blister pack per wallet and one wallet per outer carton.

The fluconazole/VIVJOA dosage regimen is in an 18-count (NDC 74695-945-18) blister package within a child resistant wallet. There is one blister pack of VIVJOA (oteseconazole capsules) per wallet and one wallet per outer carton. The outer carton and wallet contain the following: “fluconazole/VIVJOA dosage regimen” and “fluconazole is prescribed separately”.

Fluconazole is not supplied in the carton.

16.2 Storage and Handling

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F to 86°F) [See *USP Controlled Room Temperature*]. Protect from light when removed from the outer carton.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Embryo-Fetal Toxicity

Advise patients that VIVJOA is contraindicated in females of reproductive potential and in pregnant women because it may cause fetal harm [see *Contraindications (4), Warnings and Precautions (5.1) and Use in Specific Populations (8.1, 8.3)*].

Lactation

Advise patients that VIVJOA is contraindicated in lactating women because it may cause harm to the breastfed infant [see *Contraindications (4), Warnings and Precautions (5.1) and Use in Specific Populations (8.2)*].

Important Administration Instructions

Advise patients that VIVJOA must be taken with food, and that capsules must be swallowed whole and not chewed, crushed, dissolved, or opened [see *Dosage and Administration (2.1)*].

Concomitant Administration with BCRP Transporter Substrates

Advise patients to inform their health care provider if they are taking a BCRP substrate (e.g., rosuvastatin). Concomitant use with VIVJOA may increase the exposure of drugs that are BCRP substrates, which may increase the risk of adverse reactions associated with these drugs [see *Drug Interactions (7.1)*].

Manufactured for and distributed by:

Mycovia Pharmaceuticals, Inc.

Durham, NC 27703

Patent numbers: 8,236,962, 8,754,227, 10,173,998, 10,464,921, 10,836,740, 9,840,492, 10,414,751