

Section 6 updated: September 2016

## **SUMMARY OF PRODUCT CHARACTERISTICS**

Section 6 updated: September 2016

**DIETHYLCARBAMAZINE CITRATE TABLETS 100 MG USP SHOULD ONLY BE USED AS PART OF A MASS DRUG ADMINISTRATION PROGRAMME FOR ELIMINATION OF LYMPHATIC FILARIASIS IN AREAS WHERE ONCHOCERCIASIS IS NOT CO-ENDEMIC. USE AS DIRECTED.**

## **1. NAME OF THE MEDICINAL PRODUCT**

Diethylcarbamazine Citrate Tablets 100 mg USP (DEC Tablets)\*  
(diethylcarbamazine citrate)

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains 100 mg diethylcarbamazine citrate (equivalent to 50.9 mg diethylcarbamazine).

Excipient(s):

81 mg lactose monohydrate

For a full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Tablet.

White to off white, uncoated, circular, flat bevelled-edged tablets debossed with “C” on one side and plain with a break line on the other side.

The tablet can be divided into equal halves.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Diethylcarbamazine Citrate Tablets, in combination with albendazole, are indicated in adults and in children over 2 years for large scale preventative chemotherapy interventions for the control of lymphatic filariasis, following the recommendations of the WHO Global Programme to Eliminate Lymphatic Filariasis.

The mass drug administration (MDA) programme consists of administering a single annual dose of diethylcarbamazine citrate 6 mg/kg with albendazole 400 mg to affected communities, in areas where onchocerciasis is not co-endemic (see section 4.3).

### **4.2 Posology and method of administration**

Use of Diethylcarbamazine Citrate Tablets is restricted to community-wide MDA programmes in lymphatic filariasis-endemic countries following the recommendations of the WHO Global Programme to Eliminate Lymphatic Filariasis. The primary objective of the MDA strategy is to interrupt transmission of disease by reducing microfilaraemia to below 1%. In countries where onchocerciasis is not co-endemic, annual single doses each of diethylcarbamazine citrate (6 mg/kg) and albendazole (400 mg) are co-administered for 4 to 6 years.

---

\* Trade names are not prequalified by WHO. This is the responsibility of the local Drug Regulatory Authority. Throughout this WHOPAR the proprietary name is given as an example only.

Section 6 updated: September 2016

In order for MDA to be successful, at least 85% of the total population or 65% of the eligible population should receive the drugs every year for 4 to 6 years. MDA programmes that use diethylcarbamazine citrate and albendazole typically exclude children under 2 years, pregnant women and lactating mothers, the elderly, and those with chronic or debilitating illnesses. These groups account for approximately 15% of the endemic population.

Since the drugs used for MDA programmes have an established safety profile and are easy to administer, the community is not screened for the presence of microfilaraemia prior to drug administration. In view of the large number of people to be treated, MDA programmes utilize, for distribution of the drugs, the services of trained health care workers and community volunteers drawn from all sections of the society. Hence endemic communities do not usually receive drugs under direct medical supervision.

### Posology

Diethylcarbamazine Citrate Tablets should be co-administered with a single dose of albendazole 400 mg tablets (refer to prescribing information for albendazole 400 mg tablets).

#### *Adults*

The usual dose is 6 mg per kg of body weight diethylcarbamazine citrate given as a single dose administered once per year for 4 to 6 years.

#### *Children*

In children over 2 years, the usual dose is 6 mg per kg of body weight diethylcarbamazine citrate given as a single dose administered once per year for 4 to 6 years.

### Method of administration

Oral use.

Diethylcarbamazine Citrate Tablets should preferably be administered after meals.

### **4.3 Contraindications**

Co-infection with onchocerciasis. Serious ocular damage may occur.

### **4.4 Special warnings and precautions for use**

#### Use of diethylcarbamazine citrate in patients without underlying filarial infestation

Diethylcarbamazine citrate must be used with care in patients who have a history of convulsions or who show factors that predispose them to convulsions (see section 4.8).

#### *Severely ill persons*

The frail, elderly and debilitated, especially those with cardiac or renal disease, are normally excluded when diethylcarbamazine citrate is used in MDA programmes. In the event of a concomitant disorder, the patient should be allowed to recover before taking diethylcarbamazine citrate.

Diethylcarbamazine Citrate Tablets contain lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption may experience symptoms of intolerance.

Section 6 updated: September 2016

### *Children*

Children under the age of 2 years are normally excluded when diethylcarbamazine citrate is used in MDA programmes.

### Use of diethylcarbamazine citrate in patients with underlying filarial infestation

#### *Lymphatic filariasis*

Patients with symptoms suggesting lymphatic filariasis (swelling of legs, arms, female breast or male genitalia) should be managed according to the relevant local or international treatment guidelines. For WHO options for treatment, see references at end of this SmPC, under Section 4.2

#### *Loa loa*

The intensity and the severity of the undesirable effects that appear after administration of diethylcarbamazine citrate are associated with the level of microfilariae in the blood prior to treatment. In the event of *Loa loa* infestation, the level of microfilariae present in the blood is often very high, which predisposes the treated patients to an increased risk of serious side effects. Serious central nervous system problems such as encephalopathy and coma have been observed in patients with *Loa loa* infections that have been treated with diethylcarbamazine citrate, in particular.

#### *Onchocerciasis*

Diethylcarbamazine must not be administered in patients co-infected with onchocerciasis. Skin and/or systemic reactions of variable severity (Mazzotti reaction) and eye reactions have been observed after administration of drugs with rapid microfilaricidal action such as diethylcarbamazine citrate. If there is evidence of hypersensitivity to the eye, administration must be stopped due to potential sight loss. These manifestations are probably associated with an inflammatory process that is triggered following the death of microfilariae and the release of decomposition products.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Co-administration of diethylcarbamazine citrate with albendazole did not significantly alter the PK profile of either drug. Adequate absorption of both diethylcarbamazine and albendazole was observed in both adults and children aged over 5 years.

No other known interactions.

## **4.6 Pregnancy and lactation**

### Pregnancy

The potential risk to the fetus in humans is unknown.

Diethylcarbamazine Citrate Tablets should not be used in pregnancy, and as such, pregnant women are normally excluded when diethylcarbamazine citrate is used in MDA.

### Breastfeeding

It is unknown whether diethylcarbamazine or its metabolites are excreted in human milk. As a risk to the newborn or infant cannot be excluded, the product should not be given to breastfeeding women.

### Fertility

Section 6 updated: September 2016

The potential risk for humans is unknown. No specific studies with diethylcarbamazine citrate in humans have been conducted to evaluate effects on fertility.

#### **4.7 Effects on ability to drive and use machines**

Diethylcarbamazine Citrate Tablets may cause short term drowsiness which may impact on the ability to drive and use machines (see section 4.8). Vehicle drivers and machine users should be informed of the risk of drowsiness related to using this medicinal product.

#### **4.8 Undesirable effects**

<p style="text-align: center;"><b>DIETHYLCARBAMAZINE CITRATE TABLETS 100 MG USP SHOULD ONLY BE USED AS PART OF A MASS DRUG ADMINISTRATION PROGRAMME FOR ELIMINATION OF LYMPHATIC FILARIASIS IN AREAS WHERE ONCHOCERCIASIS IS NOT CO-ENDEMIC. USE AS DIRECTED.</b></p>
---

#### Summary of the safety profile

There is no clear information on the frequency of adverse reactions occurring as a result of diethylcarbamazine citrate administration. Mild to moderate adverse reactions are common, but the incidence of serious adverse reactions is considered to be very low.

In the absence of circulating microfilaraemia, the administration of diethylcarbamazine citrate, when given at the recommended dosage, may cause nausea, vomiting, abdominal pain, diarrhoea, loss of appetite, muscle pain, dizziness, drowsiness, fatigue and headache. These begin within one to two hours and may persist for several hours.

In patients with circulating microfilaraemia adverse reactions may be more common and severe, particularly in patients with a high parasite burden. Adverse reactions vary with the infecting filarial species, may be local and/or systemic, and may occur with or without fever. These are considered allergic reactions due to antigen-antibody reaction caused by dead microfilariae or adult filarial worms, and the intensity and the severity of adverse reactions are usually associated with the level of microfilariae in the blood prior to treatment. Usually such symptoms are transient and self-limiting, but when the symptoms are significant enough to interfere with daily life, the patient needs to be observed carefully and given clinically appropriate treatment. Steroids have been used. If there is evidence of hypersensitivity involving the eye, administration must be stopped due to potential sight loss (see section 4.4).

Section 6 updated: September 2016

<b>System Organ Class</b>	<b>Reactions attributable to DEC (may be seen in subjects without microfilariae)</b>	<b>Reactions attributable to death of microfilariae (may be seen in subjects with microfilariae and/or adult worm infections)</b>
Blood and lymphatic system disorders		Lymphadenitis, lymphangitis, lymph node abscess, lymph node pain, lymphoedema
Gastrointestinal disorders	Nausea, vomiting, abdominal pain, diarrhoea	Abdominal pain, nausea, vomiting, diarrhoea
General disorders and administration site conditions		Pyrexia, chills, weakness, malaise
Metabolism and nutrition disorders	Decreased appetite	Decreased appetite
Musculoskeletal and connective tissue disorders	Myalgia	Myalgia, arthralgia, chest pain
Nervous system disorders	Dizziness, somnolence, lethargy, headache	Dizziness, headache, lethargy
Renal and urinary disorders		Haematuria
Reproductive system and breast disorders		Epididymitis, spermatic cord inflammation, hydrocele, scrotal mass
Respiratory disorders		Dyspnoea, cough
Skin and subcutaneous tissue disorders		Pruritus, papular rash
Vascular disorders		Circulatory collapse, orthostatic hypotension
<b>Reactions only observed in subjects with loiasis or onchocerciasis infection (see section 4.3)</b>		
Cardiac disorders	Tachycardia	
Eye disorders	Optic neuritis, punctate keratitis, iridocyclitis, conjunctivitis, visual field defect, eye pain, lacrimation, photophobia, corneal oedema	
Immune system disorders	Mazotti reaction	
Infections and infestations	Meningoencephalitis helminthic	
Investigations	Increased intraocular pressure	
Nervous system disorders	Coma, allergic encephalitis, encephalopathy, vertigo, convulsion (isolated cases in patients with a history of epilepsy)	
Renal and urinary disorders	Proteinuria	

NOTE: frequency of all reactions is unknown (cannot be estimated from the available data).

DEC = diethylcarbamazine citrate.

### Children

Frequency, type and severity of adverse reactions in children are expected to be the same as in adults.

## **4.9 Overdose**

### Symptoms

Overdose may cause nausea, vomiting, headache, vertigo, drowsiness and, in rare but serious cases, convulsions.

### Management

Administration of activated charcoal may be of value. Monitor for adverse reactions, with symptomatic treatment and hospitalisation if necessary.

Section 6 updated: September 2016

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Anthelmintics, ATC code: P02CB02

#### Mechanism of action

Diethylcarbamazine citrate is a synthetic piperazine derivative with an anthelmintic action.

MDA of diethylcarbamazine citrate has potential to interrupt the parasitic life cycle by destruction of microfilariae which are essential for host to vector transmission of the parasite. The basic principle of MDA is to suppress microfilaraemia to levels at which transmission is not possible.

The mode of action of diethylcarbamazine citrate is not clearly known. Diethylcarbamazine citrate can be best described as a potent anti-microfilaraemic agent with variable macrofilaricidal properties. The drug is known to exert its effect directly on the parasite and also achieve parasite killing by activating the host immune response. Several potential modes of action of diethylcarbamazine citrate leading to microfilariae killing have been identified:

- Overstimulation of the neuromuscular system of the parasites and increased motility, inhibition of vital parasite metabolic enzymes, activation of surface membrane complement, activation of eosinophils and release of eosinophil-derived cationic proteins, enhanced eosinophil-dependent antibody mediated destruction of parasites and increased adhesion of parasites to phagocytic and antibody producing cells.
- Inhibition of acetylcholinesterase production by parasites, leading to increased levels of lysosomal enzymes  $\beta$ -glucuronidase and acid phosphatase that are involved in phagocytosis.
- Release of nitric oxide as evidenced by increased levels of nitrite and nitrates.
- Changes in arachidonic acid metabolism in the parasite and the host that alter adhesiveness of the parasite.

The mode of action of diethylcarbamazine citrate on adult worms is less well documented. However, there is sufficient evidence to suggest that diethylcarbamazine is macrofilaricidal. Degenerating adult worms have been demonstrated in lymph nodes post-treatment. It is also well known that the macrofilaricidal effects are inconsistent.

Diethylcarbamazine citrate was found to have effects on the arachidonic acid and cyclooxygenase metabolic pathways and COX-1 pathway of filaria. It has also been determined that inducible nitric oxide is essential for the rapid sequestration of microfilariae by diethylcarbamazine citrate. It has been suggested that as diethylcarbamazine citrate alters arachidonic acid metabolism in microfilariae (and in host endothelial cells), these changes may result in vasoconstriction and amplified endothelial adhesion, leading to immobilization of microfilarial parasites, enhanced adherence, and cytotoxic activity by host platelets and granulocytes. This would represent activation of the innate, nonspecific immune system, independent of the adaptive, antigen-specific, immune response.

#### Pharmacodynamic effects

##### *Decrease of microfilariae in blood*

Administration of diethylcarbamazine citrate (6 mg/kg) and albendazole 400 mg resulted in a decrease in blood microfilariae (See Clinical efficacy and safety section below).

Section 6 updated: September 2016

Clinical efficacy and safety

The efficacy of diethylcarbamazine citrate was first established as a 12-day treatment for lymphatic filariasis; however meta-analysis across studies has indicated that a single dose is as effective as a 12-day treatment regimen in clearance of microfilariae from the blood. Decrease in blood microfilariae is the standard measure of efficacy in lymphatic filariasis.

The improved efficacy of two-drug regimens in treating subjects with microfilariae supported the use of diethylcarbamazine citrate and albendazole in single-dose, once-yearly treatment to reduce microfilaraemia in endemic populations and this is one of the regimens used in the MDA programmes.

The effectiveness of MDA programmes has been demonstrated in studies conducted in India, Papua New Guinea, Vanuatu and Egypt. For example, the Ramaiah KD et al., 2011 study suggests that six rounds of once-yearly mass administration of single doses of diethylcarbamazine citrate and albendazole, with 60% to 70% treatment coverage, is likely to achieve total interruption of transmission and elimination of lymphatic filariasis in the majority of villages. An efficacy summary of the microfilariae clearance observed in several field studies in the MDA programmes is presented in the following table:

MDA programmes*				
Study	Treatment Arm	Number of Subjects	Microfilaraemia	
			Baseline	Endpoint
Ramaiah KD et al., 2011 Five villages (South India)	6 mg/kg DEC plus 400 mg ALB	5622 60% to 70% of the eligible population (>15 kg body weight)	Baseline Mf prevalence rate ranged from 4.27% (10/234) to 11.36% (5/44) in the 5 study villages; the overall prevalence was 8.10% (63/778) (CI: 6.18 to 10.01).	<b>After 6 rounds of MDA:</b> Overall prevalence fell to 1.01% (8/791) (CI: 0.31 to 1.71), equivalent to microfilaraemia clearance rate of 88%. The Mf prevalence fell to <0.5% in 4 of the 5 villages and it was 2.93% (6/205) in the fifth village. Two villages achieved 0% prevalence after the fourth round of MDA and maintained the same level thereafter. Reduction in Mf prevalence ( $P < 0.05$ ).

Section 6 updated: September 2016

<a href="#">Weil GJ et al., 2008</a> Four villages (Papua New Guinea)	6 mg/kg DEC plus 400 mg ALB	971 to 1000 per cycle (compliance of 72.9%)	Baseline Mf prevalence rate was 18.6%.	<b>After 3 rounds of MDA:</b> Mf clearance rates in infected persons were 76%, 95.2%, and 98.1% after 1, 2, and 3 rounds of MDA, respectively. $P = 0.013$ The difference in baseline Mf counts between the two groups was statistically significant ( $P < 0.001$ ).
<a href="#">Rajendran R et al., 2006</a> Two communities (South India)	6 mg/kg DEC plus 400 mg ALB 6 mg/kg DEC alone	DEC plus ALB: 703 to 1724 per cycle (coverage of 72.7%-94.4%; compliance of 52.4%-83.6%) DEC alone: 827 to 1432 per cycle (coverage of 67.9%-98.9%; compliance of 42.4%-80.3%)	The Mf prevalence in the 2 to 25 years age group was 5.21% in DEC plus ALB arm and 4.34% in DEC alone arm	<b>After 3 rounds of MDA:</b> The percentage reductions in Mf prevalence were higher in the DEC plus ALB arm (72.4% versus 51.4%). $P < 0.05$ in both arms.
<a href="#">Ramzy RM et al., 2006</a> Four villages (Egypt)	6 mg/kg DEC plus 400 mg ALB	Approx 1800	Microfilaraemia prevalence rates were much higher in Giza than in Qalubya (11.5 [95% CI: 10.2 to 12.8] and 3.1 [95% CI: 2.4 to 3.8], respectively).	<b>After 5 rounds of MDA:</b> In Giza, prevalence rates of microfilaraemia fell from 11.5% to 1.2%. In Qalubya, prevalence rates of microfilaraemia fell 13.6% to 3.1%. $P < 0.0001$ in both Giza and Qalubya
<a href="#">Fraser M et al., 2005</a> Eight sentinel sites (Vanuatu)	100 to 400 mg DEC (depending on age) plus 400 mg ALB	561 (average of 72% of the eligible population)	<b>Baseline:</b> Microfilaraemia prevalence was 12%	<b>After 2 rounds of treatment:</b> Prevalence of microfilaraemia was reduced by 93% (from 12% to 0.8%) Reduction in prevalence of microfilaraemia ( $P < 0.001$ )

\*Studies used various formulations of diethylcarbamazine.

## 5.2 Pharmacokinetic properties

### Absorption and bioavailability

Diethylcarbamazepine is readily absorbed following oral administration. Bioavailability is between 80 and 85%. Following single dose administration of one diethylcarbamazine tablet in healthy volunteers in the fed state, the mean ( $\pm$ SD) diethylcarbamazepine  $C_{max}$  value was 598 ( $\pm$ 84) ng/ml and the corresponding value for AUC was 7950 ( $\pm$ 1660) ng.h/ml. The mean ( $\pm$ SD) diethylcarbamazepine  $T_{max}$  value was 2.25 ( $\pm$  1.17) hours.

Section 6 updated: September 2016

Section 6 updated: September 2016

### Distribution

Diethylcarbamazine is widely distributed in tissues and is mainly excreted in the urine unchanged and as the metabolite, diethylcarbamazine N-oxide. Urinary excretion, and hence plasma half-life, is dependent on urinary pH. About 5% of a dose is excreted in the faeces.

### Metabolism

In rats and monkeys after an intravenous dose, 10-20% is excreted in the urine as unchanged drug, most of which is excreted in the first 3 hours. Metabolites more slowly eliminated include N-ethyl-4-methyl-1-piperazine-carboxamide (MEC) and their N-oxides, 4-methyl-piperazine-carboxamide and N,N-diethyl-1-piperazine-carboxamide. In vivo most of the metabolites are active on microfilariae and both N-oxides active on adults and infective larvae. The antifilarial action of DEC is swift and of short duration. This action is prolonged by the activity of metabolites, especially the N-oxides.

### Elimination

Diethylcarbamazine elimination is primarily through renal clearance, which accounts for around 50% of the total plasma clearance. The remaining clearance is via metabolism, leading to a number of metabolites, which are also renally cleared. Given that the major elimination pathway is renal clearance and that metabolic clearance is via several routes, genetic polymorphisms of drug metabolizing enzymes are very unlikely to have any clinically relevant consequences. This is borne out by the fact that there are no reported significant drug-drug interactions with diethylcarbamazine citrate which would likely be evident if modulation of metabolic clearance was an important factor in determining systemic exposure.

The PK of diethylcarbamazine citrate was not altered by timing of administration (morning versus evening).

The elimination half-life and area under the plasma concentration-time curve of diethylcarbamazine were significantly increased when an alkaline urinary pH was maintained compared with the values of these parameters obtained on a second occasion when an acidic urinary pH was maintained. This effect is unlikely to be of clinical relevance in the MDA program.

### Renal impairment

Results in patients with chronic renal impairment and in healthy subjects given a single 50 mg oral dose of diethylcarbamazine indicated that the plasma half-life of diethylcarbamazine is prolonged and its 24-hour urinary excretion is considerably reduced in those with moderate to severe renal impairment. No significant correlations were observed between age, sex or weight and renal function, but diethylcarbamazine excretion did appear to decrease with increasing urinary pH.

## **5.3 Preclinical safety data**

Nonclinical studies have not been performed with Diethylcarbamazine Citrate Tablets. However, diethylcarbamazine citrate is a well-established drug with an extensive record of clinical use.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate  
Corn starch  
Povidone (K-30)  
Magnesium stearate  
Talc

Section 6 updated: September 2016

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

60 months.

## **6.4 Special precautions for storage**

Do not store above 30°C.

Keep the bottle tightly closed in order to protect from light and moisture.

Discard the product 30 days after initial opening.

## **6.5 Nature and contents of container**

Diethylcarbamazine Citrate Tablets are supplied in bulk in a white 400 cc high density polyethylene bottle with a white polypropylene cap having an induction seal and with the inclusion of two 2 gram silica gel desiccant packets.

Each bottle contains 1000 tablets.

## **6.6 Special precautions for disposal**

Any unused product or waste material should be disposed of in accordance with local requirements.

## **7. SUPPLIER**

Eisai Co., Ltd.  
4-6-10 Koishikawa  
Bunkyo-ku  
Tokyo 112-8088  
Japan  
Phone: +81 3 3817 3684  
Fax: +81 3 3811 2710  
Email: [mailto:Eisai-asia\\_safety@hmc.eisai.co.jp](mailto:Eisai-asia_safety@hmc.eisai.co.jp)

## **8. WHO REFERENCE NUMBER (PREQUALIFICATION PROGRAMME)**

**NT002**

## **9. DATE OF FIRST PREQUALIFICATION/ RENEWAL OF THE PREQUALIFICATION**

20 August 2013

Section 6 updated: September 2016

## 10. DATE OF REVISION OF THE TEXT

August 2013. Section 6 updated in September 2016.

Detailed information on this medicinal product is available on the website of the WHO Prequalification Programme <http://www.who.int/prequal>.

### Reference List

#### **General reference:**

The major sources for the information in this SmPC are the Supatonin Tablet 50 mg Product Monograph revised in September 2008 (No. 5 edition A) and Notezine Product Monograph Sanofi-Aventis France (197497 revision 30.12.1999).

For further information, the following sources have also been used.

#### **Section 4.1 Therapeutic indications**

World Health Organization (WHO). Preparing and implementing a national plan to eliminate filariasis (in countries where onchocerciasis is not co-endemic). Geneva 2002 Ref:WHO/CDS/CPE/CEE/2000.15. Available from: [http://whqlibdoc.who.int/hq/2000/WHO\\_CDS\\_CPE\\_CEE\\_2000.15.pdf](http://whqlibdoc.who.int/hq/2000/WHO_CDS_CPE_CEE_2000.15.pdf).

World Health Organization. Preventive chemotherapy in human helminthiasis : coordinated use of anthelmintic drugs in control interventions: a manual for health professionals and programme managers. 2006. Geneva.

#### **Section 4.2 Posology and method of administration**

World Health Organization Programme to Eliminate Lymphatic Filariasis, NTD-STAG M&E Sub-working Group on Disease Specific Indicators, Meeting Report 2012  
[http://apps.who.int/iris/bitstream/10665/78611/1/WHO\\_HTM\\_NTD\\_PCT\\_2013.5\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/78611/1/WHO_HTM_NTD_PCT_2013.5_eng.pdf)

World Health Organization (WHO). Preparing and implementing a national plan to eliminate filariasis (in countries where onchocerciasis is not co-endemic). Geneva 2002 Ref:WHO/CDS/CPE/CEE/2000.15. Available from: [http://whqlibdoc.who.int/hq/2000/WHO\\_CDS\\_CPE\\_CEE\\_2000.15.pdf](http://whqlibdoc.who.int/hq/2000/WHO_CDS_CPE_CEE_2000.15.pdf).

#### **Section 4.4 Special warnings and precautions for use**

Pani S P, Subramanyam R G, Das L K, Vanamail P, Hoti S L, Ramesh J, Das P K. Tolerability and efficacy of single dose albendazole, diethylcarbamazine citrate (DEC) or co administration of albendazole with DEC in the clearance of *Wuchereria bancrofti* in asymptomatic microfilaraemic volunteers in Pondicherry, South India: A hospital based study. *Filaria Journal* 2002; 1: 11.

#### **Section 4.6 Pregnancy and lactation**

Porto I et al, Antiparasitic drugs and lactation: focus on anthelmintics, scabicides, and pediculicides. *J Hum Lact.* 2003 Nov;19(4):421-5.

#### **Section 4.8 Undesirable effects**

Bryceson AD, Warrell DA, Pope HM. Dangerous reactions to treatment of onchocerciasis with diethylcarbamazine. *Br Med J.* 1977 Mar;1(6063):742-4.

Dreyer G, Pires ML, de Andrade LD, Lopes E, Medeiros Z, Tenorio J, et al. Tolerance of diethylcarbamazine by microfilaraemic and amicrofilaraemic individuals in an endemic area of Bancroftian filariasis, Recife, Brazil. *Trans R Soc Trop Med Hyg.* 1994 Mar-Apr;88(2):232-6.

Hawking F. Diethylcarbamazine and new compounds for the treatment of filariasis. *Adv Pharmacol Chemother.* 1979;16:129-94.

Horton J, Witt C, Ottesen E A, Lazdins J K, Addiss D G, Awadzi K, Beach M J, Belizario V Y, Dunyo S K, Espinel M, Gyapong J O, Hossain M, Ismail M M, Jayakody R L, Lammie P J, Makunde W, Richard L D, Selve B, Shenoy R K, Simonsen P E, Wamae C N, Weerasooriya M V. An analysis of the safety of the single

Section 6 updated: September 2016

dose, two drug regimens used in programmes to eliminate lymphatic filariasis. *Parasitology* 2000; 121 suppl: S147–60.

MacKenzie CD, Kron, M. A. Diethylcarbamazine: Review of pharmacology, mechanisms of action and clinical uses: *Trop. Dis. Bull.* 82(10), R1-R37 (1985).

McLaughlin S I, Radday J, Michel M C, Addiss D G, Beach M J, Lammie P J, Lammie J, Rheingans R, Lafontant J. Frequency, severity, and costs of adverse reactions following mass treatment for lymphatic filariasis using diethylcarbamazine and albendazole in Iogane, Haiti, 2000. *American Journal of Tropical Medicine and Hygiene* 2003; 68 (5): 568–73.

Ottesen EA. Description, mechanism and control of reactions to treatment in the human filariases. *Filariasis. Ciba Found Symp* 127. Chichester: Wiley; 1987. p. 265-283.

WHO. Weekly epidemiological record (WER). 2011 Aug 26;86(35):377–388. Available from: <http://www.who.int/wer>.

#### **Section 4.9 Overdose**

Orisakwe OE et al. Acceleration of body clearance of diethylCarbamazine by oral activated charcoal. *Pharmacol Res.* 2000 Aug;42(2):167-70.

#### **Section 5.1 Pharmacodynamic properties**

Bhattacharya C, Singh RN, Nisra S, Rathaur S. Diethylcarbamazine effect on lysosomal enzymes and acetylcholine in *Wucheria Bancroftian* infection. *Trop. Med. Int. Health.* 2, 686-690 (1997)

Cao WC, Van der Ploeg CPB, Plaisier AP, Sivera van der Sluijs J, Habbema JDF. Ivermectin for the chemotherapy of bancroftianfilariasis: a meta-analysis of the effect of single treatment. *Trop Med Int Health.* 1997 Apr;2(4):393-403.

Eberhard ML, Hightower AW, Addis DG, Lammie MJ. Clearance of *Wucheria bancrofti* antigen after treatment with diethylcarbamazine or ivermectin. *Am. J. Trop. Med. Hyg.* 57, 483-486 (1997).

Figueredo-Silva J, Jungmann P, Norões J, Piessens WF, Coutinho A, Brito C, Rocha A, Dreyer G. Histological evidence for adulticidal effect of low doses of diethylcarbamazine in bancroftian filariasis. *Trans R Soc Trop Med Hyg.* 1996 Mar-Apr; 90(2):192-4.

Fraser M, Taleo G, Taleo F, Yaviong J, Amos M, Babu M, et al. Evaluation of the program to eliminate lymphatic filariasis in Vanuatu following two years of mass drug administration implementation: results and methodologic approach. *Am J Trop Med Hyg.* 2005 Oct;73(4):753-8.

Gyapong JO, Kumaraswami V, Biswas G, Ottesen EA. Treatment strategies underpinning the global programme to eliminate lymphatic filariasis. *Expert Opin Pharmacother.* 2005 Feb;6(2):179-200.

Hoti S L, Pani S P, Vanamail P, Athisaya M K, Das L K, Das P K. Effect of a single dose of diethylcarbamazine, albendazole or both on the clearance of *Wuchereria bancrofti* microfilariae and antigenaemia among microfilaria carriers: a randomized trial. *National Medical Journal of India* 2010; 23 (2): 72–6.

Horton J, Witt C, Ottesen E A, Lazdins J K, Addiss D G, Awadzi K, Beach M J, Belizario V Y, Dunyo S K, Espinel M, Gyapong J O, Hossain M, Ismail M M, Jayakody R L, Lammie P J, Makunde W, Richard L D, Selve B, Shenoy R K, Simonsen P E, Wamae C N, Weerasooriya M V. An analysis of the safety of the single dose, two drug regimens used in programmes to eliminate lymphatic filariasis. *Parasitology* 2000; 121 suppl: S147–60.

Ismail MM, Weil GJ, Jayahinghe KS et al. Prolonged clearance of microfilaraemia in patients with Bancroftian filariasis after multiple high doses of ivermectin or diethylcarbamazine. *Trans. R. Soc. Trop. Med. Hyg.* 90, 684-688 (1996).

Kshirsagar NA, Gogtay NJ, Garg BS, Deshmukh PR, Rajgor DD, Kadam VS, et al. Safety, tolerability, efficacy and plasma concentrations of diethylcarbamazine and albendazole co-administration in a field study in an area endemic for lymphatic filariasis in India. *Trans R Soc Trop Med Hyg.* 2004 Apr;98(4):205-17.

McGarry HF, Plant LD, Taylor MJ. Diethylcarbamazine activity against *Brugiamalayi* microfilariae is dependent on inducible nitric-oxide synthase and the cyclooxygenase pathway. *Filaria J.* 2005 Jun 2;4:4.

MacKenzie CD, Kron, M. A. Diethylcarbamazine: Review of pharmacology, mechanisms of action and clinical uses: *Trop. Dis. Bull.* 82(10), R1-R37 (1985).

Section 6 updated: September 2016

McLaughlin S I, Radday J, Michel M C, Addiss D G, Beach M J, Lammie P J, Lammie J, Rheingans R, Lafontant J. Frequency, severity, and costs of adverse reactions following mass treatment for lymphatic filariasis using diethylcarbamazine and albendazole in Iogane, Haiti, 2000. *American Journal of Tropical Medicine and Hygiene* 2003; 68 (5): 568–73.

Maizels RM, Denham DA. Diethylcarbamazine (DEC): immunopharmacological interactions of an anti-filarial drug. *Parasitology*. 1992;105 Suppl:S49-60.

Melrose WD. Chemotherapy for lymphatic filariasis: progress but not perfection. *Expert Rev Anti Infect Ther*. 2003 Dec;1(4):571-7.

Ottesen EA. Efficacy of diethylcarbamazine in eradicating infection with lymphatic-dwelling filariae in humans. *Rev Infect Dis*. 1985 May-Jun;7 (3):341-56.

Pani S P, Subramanyam R G, Das L K, Vanamail P, Hoti S L, Ramesh J, Das P K. Tolerability and efficacy of single dose albendazole, diethylcarbamazine citrate (DEC) or co administration of albendazole with DEC in the clearance of *Wuchereria bancrofti* in asymptomatic microfilaraemic volunteers in Pondicherry, South India: A hospital based study. *Filaria Journal* 2002; 1: 11.

Rajendran R, Sunish IP, Mani TR, Munirathinam A, Arunachalam N, Satyanarayana K, Dash AP. Community-based study to assess the efficacy of DEC plus ALB against DEC alone on bancroftian filarial infection in endemic areas in Tamil Nadu, south India. *Trop Med Int Health*. 2006 Jun;11(6):851-61.

Ramaiah KD, Vanamail P, Yuvaraj J, Das PK. Effect of annual mass administration of diethylcarbamazine and albendazole on bancroftian filariasis in five villages in south India. *Trans R Soc Trop Med Hyg*. 2011 Aug;105(8):431-7.

Ramzy RM, El Setouhy M, Helmy H, Ahmed ES, Abd Elaziz KM, Farid HA, et al. Effect of yearly mass drug administration with diethylcarbamazine and albendazole on bancroftian filariasis in Egypt: a comprehensive assessment. *Lancet*. 2006 Mar 25;367(9515):992—9.

Reddy G S, Vengatesvarlou N, Das P K, Vanamail P, Vijayan A P, Kala S, Pani S P. Tolerability and efficacy of single-dose diethylcarbamazine (DEC) or ivermectin in the clearance of *Wuchereria bancrofti* microfilaraemia in Pondicherry, South India. *Tropical Medicine and International Health* 2000; 5 (11): 779–85.

Sean C Sweetman (Editor) *Anthelmintics in Martindale The Complete Drug Reference* 36th Edition 2009; 143-4.

Taylor MJ, Cross HF, Mohammed AA, Trees AJ, Bianco AE. Susceptibility of *Brugia malayi* and *Oncocerca lienalis* microfilariae to nitric oxide and hydrogen peroxide in cell free culture and from IFN  $\gamma$ -activated macrophages. *Parasitol*. 112, 315-322 (1996).

Weil GJ, Sethumadhavan KVP, Santhanam S, Jain DC, Ghosh TK. Persistence of parasite antigenaemia following diethylcarbamazine therapy of Bancroftian filariasis. *Am. J. Trop. Med. Hyg*. 59, 523-525 (1988).

Weil GJ, Kastens W, Susapu M, Laney SJ, Williams SA, King CL, et al. The impact of repeated rounds of mass drug administration with diethylcarbamazine plus albendazole on bancroftian filariasis in Papua New Guinea. *PLoS Negl Trop Dis*. 2008;2(12):e344.

Winkler S, EL Menyawi, Floris-Linnau K, Graninger W. Short report: total serum levels of the nitric oxide derivatives nitrite/nitrate during microfilarial clearance in human filarial disease. *Am. J. Trop. Med. Hyg*. 59, 523-525 (1998).

### Section 5.2 Pharmacokinetic properties

Bangham DR. Metabolism and excretion of <sup>14</sup>C-labelled diethylcarbamazine. *Br J Pharmacol*. 1955;10:397-405.

Edwards G, Breckenridge AM, Adjepon-Yamoah KK, Orme M L'E, Ward SA. The effect of variations in urinary pH on the pharmacokinetics of diethylcarbamazine. *Br J Clin Pharmacol* (1981) 12, 807–12.

Ilondu N, Orisakwe O E, Ofoefule S, Afonne O J, Obi E, Chilaka K C, Orish C. Pharmacokinetics of diethylcarbamazine: Prediction by concentration in saliva. *Biological and Pharmaceutical Bulletin* 2000; 23 (4): 443–5.

Sweetman SC, editor. *Anthelmintics*. In: *Martindale: The Complete Drug Reference*. 35th ed. London: Pharmaceutical Press; 2007. p. 125-6.