



TAMPER EVIDENT: Do not use if the neckband printed with [ ] is broken or missing.

Code 128  
70236M

2 3/4" 1 11/16"

**Drug Facts**

**Active ingredient (in each 5 mL)** Purpose  
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide..... Cough suppressant

**Uses** temporarily relieves  
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants  
 ■ the impulse to cough to help you get to sleep

**Warnings**  
**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have**  
 ■ chronic cough that lasts as occurs with smoking, asthma or emphysema  
 ■ cough that occurs with too much phlegm (mucus)  
**Stop use and ask a doctor** if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.  
**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ shake bottle well before use  
 ■ measure only with dosing cup provided  
 ■ do not use dosing cup with other products  
 ■ dose as follows or as directed by a doctor  
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

**Other information** ■ each 5 mL contains: sodium 7 mg  
 ■ store at 20-25°C (68-77°F) ■ dosing cup provided

**Inactive ingredients** citric acid, edetate disodium, ethylcellulose, FD&C Yellow No. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

**Questions? 1-888-963-3382**  
 You may also report side effects to this phone number.

**Reckitt Benckiser**  
 C I A 70236M  
 PT# C I A 60333  
 8137779  
 Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224 ©2014 RB 022114

**12 Hour Cough Relief**  
**Children's Delsym**  
 Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

**12 Hour Cough Relief**  
**Children's Delsym**  
**COUGH**  
**Orange**  
 Flavored Liquid

NDC 63824-177-65  
**Children's Delsym**  
 Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

**12 Hour Cough Relief**  
**Children's Delsym**  
 Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant)

**COUGH**

**COUGH**

**COUGH**

**Also Available In Grape Flavor**



Age 4+

For Children & Adults



Please visit our web site:  
[www.delsym.com](http://www.delsym.com)

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Orange**  
Flavored Liquid 148 mL (5 fl oz)  
Day or Night Alcohol-free

**DELSYM® DOSING**  
 SHAKE WELL BEFORE USE.  
 Measure only with dosing cup provided. Do not use dosing cup with other products. mL = milliliter

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.



**Dosing Cup Included**

**Contains No Fever Reducer or Pain Reliever**

3-63824-27665-6  
(UPC @ 100% truncated)

Lot No.:  
Exp. Date:

5 11/16"

