

Anaroxyl[®]

- aqueous solution
for s.c. i.m. or i.v. use
- tablet for oral use

Composition

Each ml of the aqueous solution contains: 5 mg Monosemicarbazone Adrenochrome 2 aq.

Each tablet contains: 2.5 mg Monosemicarbazone Adrenochrome 2 aq.

Characteristics

Anaroxyl[®] is a haemostatic with a rapid onset of action. It reduces normal and pathologic bleeding time by decreasing capillary permeability.

Indications

Prevention and treatment of surgical and non-surgical capillary bleedings.

Dosage

For the prevention and treatment of capillary bleedings, accordingly to the severity of the case, 4-12 tablets divided over the day, preferably one hour before meals. Prior to surgery, medication is started on the preceding evening, either orally or by intramuscular or subcutaneous injection of usually 1.5-5 mg. In addition, an injection is given about half an hour before the surgical intervention.

Administration

Anaroxyl[®] solution should be administered by intramuscular or subcutaneous injection; in emergency cases the injections may be given intravenously. The tablets should be taken orally.

Contra-indications

There are no known contra-indications.

Use in Pregnancy

During pregnancy, the benefits of the use of **Anaroxyl[®]** should be weighed against the possible hazards of the foetus.

Warnings and precautions

Because of the content of diprophylline (90mg/ml), the dosage of the 5mg/ml injection preparation to premature babies, infants and young children should not exceed 0.3 ml.

Adverse reaction

No adverse reactions have been reported at the recommended dosages.

Storage Condition

Store in a cool (in between 2°-25° C), dry place & protect from light. Keep out of the reach of children.

Commercial Pack

Anaroxyl[®] injection: Each box contains 5x1 ml ampoules in Alu-PVC blister strip.

Anaroxyl[®] tablet: Each box contains 3x10 tablets in Alu-Alu blister strip.



Manufactured by:
Nuvista Pharma Limited
Tongi - Gazipur